arXiv:1609.03638v1 [math.AG] 13 Sep 2016

MONODROMY AND ALGEBRAIC CYCLES

KALYAN BANERJEE

Abstract. In this text we are going to discuss the relation between monodromy and algebraic cycles.

1. Introduction

This article is motivated by the following question due to Claire Voisin in [Vo]. Let S be a smooth projective complex algebraic embedded in a projective space $\mathbb{P}^N$. Let H_t be a hyperplane in $\mathbb{P}^N$ such that $S \cap H_t = C_t$ is smooth and irreducible. Let us consider the closed embedding j_t from C_t to S and consider the push-forward homomorphism j_{t*} from $A_0(C_t)$ to $A_0(S)$, where A_0 denotes the group of algebraically trivial algebraic cycles modulo rational equivalence. Then the natural question is that what is the kernel of j_{t*}. In [BG] it was proved that this kernel is a countable union of translates of an abelian subvariety of $J(C_t)$ (as conjectured by Voisin in [Vo]). Furthermore we can prove that for a very general hyperplane section C_t of S, this abelian variety is either trivial or all of $J(C_t)$, in the case when S is a K3 surface. This was done using an elegant technique in [Voi] concerning monodromy of a Lefschetz pencil of hyperplane sections on S. In [BG] we have further generalised the monodromy technique present in [Voi] using étale fundamental group and étale monodromy over an arbitrary ground field which is uncountable and of characteristic zero. Furthermore if $A_0(S)$ is not isomorphic to $Alb(S)$, which is the case if S is K3, then for a very general hyperplane section the kernel of j_{t*} is actually countable.

The aim of this paper is to address the following two questions. One is consider a fibration on a K3 surface (over an uncountable ground field of characteristic zero) which is close to being a Lefschetz pencil. That

[0]Mathematics Classification Number: 14C25, 14C21, 14C30, 14D05, 14D20, 14D21

[0]Keywords: Pushforward homomorphism, K3 surfaces, Monodromy, Chow groups.

is a morphism from $S \to C$ such that there exists some Lefschetz pencil $S \to D$ and we have the obvious triangle commutative. Then what can we say about the kernel of the push-forward homomorphism for a very general t in C. First we prove that for any t such that C_t is smooth and irreducible curve the kernel of the push-forward is a countable union of translates of an abelian subvariety A_t of $J(C_t)$, where S_t is fiber over t and is a smooth irreducible curve. Then by appealing to the étale monodromy argument we get that this abelian variety A_t is either zero or $J(C_t)$ for a very general t. This is done by reducing everything to the monodromy of the Lefschetz pencil $S \to D$. Furthermore following techniques in [BG] we get that this kernel is actually countable for a very general t in C. So we get the following theorem.

For a fibration $S \to C$, which is "close to be a Lefschetz pencil", for a very general $t \in C$ we have that the kernel of j_{t*} is countable.

The second question is to address the understanding variance of Voisin's question in the case of considering the natural homomorphism from $A_0(C_t) \times A_0(C_s) \to A_0(S)$ (S a K3 surface over the field of complex numbers), when we consider a net $S \to \mathbb{P}^1 \times \mathbb{P}^1$, such that each pencil in this net is a Lefschetz pencil. Then by using the monodromy technique we prove that for a very general t, s we have the abelian subvariety $A_{(t,s)}$ of $J(C_t) \times J(C_s)$ is either zero or $J(C_t)$ or $J(C_s)$ or $J(C_t) \times J(C_s)$. Since S is K3 surface we can rule out the possibility that $A_{(t,s)}$ is equal to $J(C_t) \times J(C_s)$. So we get the following result:

For a very general t, s in the net, we have that the kernel of $j_{t,s*}$ from $A_0(C_t) \times A_0(C_s)$ to $A_0(S)$ is countable.

Also we consider the case when we consider a Lefschetz pencil on S a K3 surface and consider $C_t \times C_t$ inside $S \times S$. Then we prove that:

For a very general t the kernel of j_{t*} from $A_0(C_t \times C_t)$ to $A_0(S \times S)$ is either zero or isogenous to $J(C_t)$ or is $Alb(C_t \times C_t)$.

The other one is to understand the Branched covers of a $K3$ surface and curves on that. Let $\widetilde{S} \to S$ be a branched cover of a K3 surface. Then consider a Lefschetz pencil on S, then for a general member C_t of the pencil, $\widetilde{C}_t$ (the preimage of C_t) is smooth and irreducible. Then we address the question of theh kernel of j_{t*} from $A_0(\widetilde{C}_t)$ to $A_0(S)$. We prove the following result.

The kernel of j_{t*} is a countable union of translates of an abelian subvariety A_t in $J(\widetilde{C}_t)$. For a general t this abelian variety A_t contains an abelian subvariety B_t, which is either zero or isogenous to $J(C_t)$. Furthermore B_t is actually zero for a very general t.

The organisation of the text is as follows. In the second section we discuss the behaviour of monodormy for an arbitrary fibration which is close to being a Lefschetz pencil. In the third section we discuss the relation between monodromy and algebraic cycles on various constructions arising from a K3 surface and taking a Lefschetz pencil or a net on it.

Acknowledgements: The author would like to thank the ISF-UGC project hosted by Indian Statistical Institute, Bangalore center.

2. Fibration over a smooth projective curve and monodromy

Let S be an even dimensional smooth projective surface with irregularity zero and suppose that we have a regular flat projective morphism f from S to a smooth projective curve C. Let t be a point on C such that the scheme theoretic fiber over t, that is C_t is non-singular. We consider the closed embedding j_t of C_t into S. Then we have j_{t*} from $A^1(C_t)$ to $A^2(S)$. We want to understand the kernel of this push-forward homomorphism j_{t*}. We have $A^1(C_t)$ is regularly isomorphic to the Jacobian variety $J(C_t)$. Then by [BG] proposition 6 we have the following theorem.

Theorem 2.1. *The kernel of the push-forward homomorphism j_{t*} from $J(C_t)$ to $A^2(S)$ is a countable union of translates of an abelian subvariety A_t of $J(C_t)$.*

Proof. See proposition 6 in [BG]. □

Now consider an embedding of S into a projective space $\mathbb{P}^N$, consider a line D in $\mathbb{P}^{N*}$, such that it gives a Lefschetz pencil on S. Then since C is a smooth projective curve there is a finite map π from C to this line D. Suppose that the composition $S \to \mathbb{P}^1$ is same as $S \to C \to \mathbb{P}^1$. Then except for the branch locus of π, we have a finite covering from U to U', where U, U' are open sets of C, D respectively, we get that for a point t in U, C_t is isomorphic to $D_{\pi(t)}$, where $D_{\pi(t)}$ denotes the

fiber over the point $\pi(t)$ in D. Shrinking U, U' further we can assume that $C_t, D_{\pi(t)}$ are smooth. This isomorphism induces an isomorphism of $J(C_t)$ and $J(D_{\pi(t)})$. Call that isomorphism η_t. Then we have the following commutativity at the level of Chow groups.

$$\begin{array}{ccc} J(C_t) & \xrightarrow{j_{t*}} & A^2(S) \\ \downarrow{\scriptstyle \eta_t} & & \downarrow{\scriptstyle \mathrm{id}} \\ J(D_{\pi(t)}) & \xrightarrow{i_{t*}} & A^2(S) \end{array}$$

This gives us that

$$\eta_t(\ker(j_{t*})) = \ker(i_{t*}) \ .$$

By the previous theorem we get that $\ker(j_{t*})$ is a countable union of translates of an abelian variety A_t, and similarly $\ker(i_{t*})$ is a countable union of translates of an abelian variety B_t. So write

$$\ker(j_{t*}) = \cup_{i\in\mathbb{N}} x_i + A_t$$

therefore

$$\eta_t(\ker(j_{t*})) = \cup_{i\in\mathbb{N}} \eta_t(x_i) + \eta_t(A_t)$$

Since $\eta_t(\ker(j_{t*}))$ is $\ker(i_{t*})$ we get that

$$\cup_{j\in\mathbb{N}} y_j + B_t = \cup_{i\in\mathbb{N}} \eta_t(x_i) + \eta_t(A_t) \ .$$

Assuming that the ground field k is uncountable we get that any projective variety cannot be written as a countable union of proper Zariski closed subsets. From this it follows that

$$\eta_t(A_t) = B_t \ .$$

Now by the monodromy argument as present in [BG] we get that B_t is zero or $J(D_{\pi(t)})$ for a very general t, this gives us that $\eta_t(A_t)$ is either zero or $J(D_{\pi(t)}$, from which it follows that A_t is zero or $J(C_t)$, since η_t is an isomorphism of abelian varieties.

We can also prove that A_t is zero or $J(C_t)$ without using the isomorphism η_t and using the very nice monodromy argument and the spread argument present in [BG]. We present it here.

Theorem 2.2. *For a very general t in C, the abelian variety A_t is either zero or all of $J(C_t)$.*

Proof. Let $\bar{\eta}$ be the geometric generic point of C. Then the Jacobian of $C_{\bar{\eta}}$ is isomorphic to C_t for a very general t. So if we consider the push-forward homomorphism $j_{\bar{\eta}*}$ from $J(C_{\bar{\eta}})$ to $A^2(S_{\bar{\eta}})$ we get that the kernel of this homomorphism is a countable union of translates of an abeliab subvariety $A_{\bar{\eta}}$ of $J(C_{\bar{\eta}})$. Let L be a finitely generated extension of $k(C)$ in $k(\bar{C})$, such that $A_{\bar{\eta}}$ and $J(C_{\bar{\eta}})$ are defined over L. Let C' be a curve such that $k(C') = L$ and we have a finite morphism from C' to C. Then over a Zariski open subset U' of C' we spread $A_{\bar{\eta}}$ and $J(C_{\bar{\eta}})$, to get abelian schemes $\mathscr{A}$, $\mathscr{J}$ over U'. Let α, β be the morphisms from $\mathscr{A}$, $\mathscr{J}$ to U'. Then consider the constant sheaf $\mathbb{Z}/l^n\mathbb{Z}$ on $\mathscr{A}$, $\mathscr{J}$. Throwing out some more points from U' we can assume that α, β are non-singular, that is the fibers are non-singular, also α, β are locally projective therefore they are proper. So the higher direct image sheaves $R^1\alpha_*(\mathbb{Z}/l^n\mathbb{Z})_{\mathscr{A}}$, $R^1\beta_*(\mathbb{Z}/l^n\mathbb{Z})_{\mathscr{J}}$ are locally constant sheaves on U. Since there is an equivalence of locally constant sheaves on U and $\pi_1(U, \bar{\eta})$ modules. We get that the stalks of the above sheaves at the point $\bar{\eta}$ are $\pi_1(U, \bar{\eta})$ modules. By the proper base change theorem we have $(R^1\alpha_*(\mathbb{Z}/l^n\mathbb{Z})_{\mathscr{A}})_{\bar{\eta}}$ is isomorphic to $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Z}/l^n\mathbb{Z})$ and $(R^1\beta_*(\mathbb{Z}/l^n\mathbb{Z})_{\mathscr{J}})_{\bar{\eta}}$ is isomorphic to $H^1_{\acute{e}t}(J(C_{\bar{\eta}}), \mathbb{Z}/l^n\mathbb{Z})$ and they are $\pi_1(U, \bar{\eta})$ modules. By taking the inverse limit of this cohomologies we get that $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Q}_l)$ and $H^1_{\acute{e}t}(J(C_{\bar{\eta}}), \mathbb{Q}_l)$ are $\pi_1(U, \bar{\eta})$ modules. Also the natural morphism from $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Q}_l)$ to $H^1_{\acute{e}t}(J(C_{\bar{\eta}}), \mathbb{Q}_l)$ is a map of $\pi_1(U, \bar{\eta})$ modules as it is induced from the regular morphism $\mathscr{A} \to \mathscr{J}$.

Since we have a finite map from C to $D \cong \mathbb{P}^1$, we have a map from $\overline{k(D)}$ to $\overline{k(C)}$, which gives us a morphism of schemes $\mathrm{Spec}(\overline{k(C)})$ to $\mathrm{Spec}(\overline{k(D)})$. $\mathrm{Spec}(\overline{k(C)})$ is nothing but $\bar{\eta}$ and denote $\mathrm{Spec}(\overline{k(D)})$ as $\bar{\xi}$. Then $\bar{\eta}$ maps to $\bar{\xi}$. So consider the following fiber square

$$\begin{array}{ccc} C \times_D \bar{\xi} & \longrightarrow & \bar{\xi} \\ \downarrow & & \downarrow \\ C & \xrightarrow{i_{t*}} & D \end{array}$$

Since $\bar{\eta}$ maps to $\bar{\xi}$ we have that $\bar{\eta}$ is in $C \times_D \bar{\xi}$. Since k is uncountable we can always choose t such that we have that C_t is isomorphic to $C_{\bar{\eta}}$ and $C_t \cong D_{\pi(t)}$, and also $D_{\pi(t)}$ is isomorphic to $D_{\bar{\xi}}$. Therefore we get

that $C_{\bar{\eta}}$ is isomorphic to $D_{\bar{\xi}}$ as schemes over Spec($\mathbb{Q}$) but may not be over Spec(k). Therefore we have that $H^1_{\acute{e}t}(J(C_{\bar{\eta}}), \mathbb{Q}_l)$ is isomorphic to $H^1_{\acute{e}t}(J(D_{\bar{\xi}}), \mathbb{Q}_l)$. Now $H^1_{\acute{e}t}(J(D_{\bar{\xi}}), \mathbb{Q}_l)$ is isomorphic to $H^1_{\acute{e}t}(D_{\bar{\xi}}, \mathbb{Q}_l)$ and the map is a map of $\pi_1(U', \bar{\xi})$ modules, where U' is such that $\pi^{-1}(U') = U$. Therefore as in section 4 of [BG] we get that $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Q}_l)$ is included in $H^1_{\acute{e}t}(J(C_{\bar{\eta}}), \mathbb{Q}_l)$ which is isomorphic to $H^1_{\acute{e}t}(J(D_{\bar{\xi}}), \mathbb{Q}_l)$ and that is again isomorphic to $H^1_{\acute{e}t}(D_{\bar{\xi}}, \mathbb{Q}_l)$ on which we have the irreducibility of the tame fundamental group $\pi_1^{tame}(U', \bar{\xi})$ given by the Picard Lefshcetz formula. By the Picard Lefschetz formula and the fact that $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Q}_l)$ is a $\pi_1(U, \bar{\eta})$ module whose image is a finite index subgroup in $\pi_1(U', \bar{\xi})$, it follows that $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Q}_l)$ is $\pi_1^{\text{tame}}(U', \bar{\xi})$ equivariant. Hence we get that $H^1_{\acute{e}t}(A_{\bar{\eta}}, \mathbb{Q}_l)$ is either zero or $H^1_{\acute{e}t}(J(C_{\bar{\eta}}), \mathbb{Q}_l)$. Therefore it follows that $A_{\bar{\eta}}$ is either zero or $J(C_{\bar{\eta}})$ by Tate module reasons. Since $J(C_{\bar{\eta}}), A_{\bar{\eta}}$ are isomorphic to $J(C_t), A_t$ for a very general t, it follows that for a very general t, A_t is either zero or $J(C_t)$. Also observe that if $A_{\bar{\eta}}$ is zero then A_t is zero for all t belonging to the complement of countable union of points in C and if $A_{\bar{\eta}}$ is $J(C_{\bar{\eta}})$ then $A_t = J(C_t)$ for all t belonging to the complement of countable union of points in C. □

Lemma 2.3. *The set $C^\sharp$ consisting of all points t such that $A_t = J(C_t)$ is constructible.*

Proof. Proof goes in the same line as in lemma 17 in [BG]. □

Theorem 2.4. *Suppose that $A^2(S)$ is not weakly rationally representable. Then for a very general t in C we have the kernel of j_{t*} to be countable.*

Proof. First we show by using the constructibility of the above set $C^\sharp$ that if j_{t*} is zero for a very general t then actually j_{t*} is zero for a general t. Then we take $C_{\bar{\eta}}$, since $A^2(C_{\bar{\eta}}) = 0$ we get that there exists a curve $\Gamma_{\bar{\eta}}$, a correspondence $Z_{\bar{\eta}}$ on $\Gamma_{\bar{\eta}} \times C_{\bar{\eta}}$ such that $Z_{\bar{\eta}*}$ is onto from $J(\Gamma_{\bar{\eta}})$ to $A^1(C_{\bar{\eta}})$. We can very well choose this curve $\Gamma_{\bar{\eta}}$ to $C_{\bar{\eta}}$, and $Z_{\bar{\eta}}$ to be the diagonal in the two fold product of $C_{\bar{\eta}}$ and then we spread $C_{\bar{\eta}}$ to a surface S' and the diagonal to a correspondence $\mathscr{Z}$ over a Zariski open subset V of some curve C', such that $C' \to C$ we have a finite map. Then arguing as in theorem 19 in [BG] we prove that $A^2(S')$ tensored with $\mathbb{Q}$ is equal to the direct sum of the image of $\mathscr{Z}_*$ and images of the homomorphisms j_{t*}. Then by using the fact that $j_{t*} = 0$ for all but a finitely many t, we prove that the above direct sum

is finite and each of the summands is weakly representable, so gives rational weak representability of $A^2(S')$. Since S' is a blow up of S, it follows that $A^2(S)$ is rationally weakly representable contradicting our assumption.

All this arguments are taken from [BG] theorem 19, where they first appeared. □

3. Curves on a surface and monodromy

Let S be a smooth, projective, surface over $\mathbb{C}$. Let us fix an embedding of S inside $\mathbb{P}^N$. Let t be a closed point in $\mathbb{P}^{N*}$, consider the corresponding hyperplane H_t inside $\mathbb{P}^N$ and consider its intersection with S, then we get a curve C_t inside S. By Bertini's theorem, a general such hyperplane section of S will be smooth and irreducible. Now consider two such curves C_t, C_s in S. Then we have the following commutative diagram.

$$\begin{array}{ccc} \mathrm{Sym}^n C_t \times \mathrm{Sym}^n C_s & \longrightarrow & \mathrm{Sym}^{2n} S \\ \downarrow & & \downarrow \\ A_0(C_t) \times A_0(C_s) & \longrightarrow & A_0(S) \end{array}$$

Here the morphism from $\mathrm{Sym}^n C_t \times \mathrm{Sym}^n C_s$ to $\mathrm{Sym}^{2n} S$ is given by

$$(\sum_i P_i, \sum_j Q_j) = \sum_i P_i + \sum_j Q_j$$

and the homomorphism from $A_0(C_t) \times A_0(C_s)$ to $A_0(S)$ is given by

$$j_{t,s*} = j_{t*} + j_{s*} \ .$$

It is easy to see that the above diagram is commutative (since $\mathbb{C}$ is algebraically closed). By the Abel-Jacobi theorem $A_0(C_t) \times A_0(C_s)$ is isomorphic to $J(C_t) \times J(C_s)$. Following the argument of [BG], proposition 6 we get that the kernel of $j_{t,s*}$ is a countable union of translates of an abelian subvariety of $J(C_t) \times J(C_s)$. Call this abelian subvariety $A_{(t,s)}$. Now consider a net on S, that is two Lefschetz pencils D_1, D_2 on S. Then for a general (t, s) on $D_1 \times D_2$, the curves C_t, C_s will be smooth and irreducible. Now we prove that for a general (t, s), $A_{(t,s)}$

will either be $\{0\}$ or $J(C_t)$ or $J(C_s)$ or all of $J(C_t) \times J(C_s)$. Suppose also that $H^3(S, \mathbb{Q}) = 0$. This is the case for example of a K3 surface.

Theorem 3.1. *For a very general (t, s) in $D_1 \times D_2$, the abelian variety $A_{t,s}$ is either $\{0\}$ or $J(C_t)$ or $J(C_s)$ or $J(C_t) \times J(C_s)$.*

Proof. The argument comes from monodromy. We have a natural monodromy representation of the fundamental group of $D_1 \setminus \{0_1, \cdots, 0_m\}$ and $D_2 \setminus \{0'_1, \cdots, 0'_n\}$ on the Gysin kernels $H^1(C_t, \mathbb{Q})$ and $H^1(C_s, \mathbb{Q})$ respectively, for a very general t, s such that C_t, C_s are smooth. By theorem 3.27 in [Vo] we have that these monodromy representations are irreducible. So it will follow that the induced representation of $G = \pi_1(D_1 \setminus \{0_1, \cdots, 0_m\}, t) \times \pi_1(D_1 \setminus \{0'_1, \cdots, 0'_n\}, s)$ on $H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$ has the following property. Any G invariant subspace of it is either $\{0\}$ or $H^1(C_t, \mathbb{Q})$ or $H^1(C_s, \mathbb{Q})$ or all of $H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$. Consequently, by using the correspondence between Hodge structures of weight one and abelian varieties we have that the only non-trivial abelian subvarieties of $J(C_t) \times J(C_s)$ are either $J(C_t)$ or $J(C_s)$. Now to prove that $A_{(t,s)}$ is either one of these four possibilities we have to show that the Hodge structure corresponding to $A_{(t,s)}$ is G equivariant. So for a general t, s we have an abelian subvariety $A_{(t,s)}$ of $J(C_t) \times J(C_s)$. Now consider the isomorphism of $\mathbb{C}$ with $\overline{\mathbb{C}(t, s)}$ and view $A_{(t,s)}$ and $J(C_t) \times J(C_s)$ as abelian varieties over $\overline{\mathbb{C}(t, s)}$. Let L be the minimal field of definition of $A_{(t,s)}$ and $J(C_t) \times J(C_s)$ in $\overline{\mathbb{C}(t, s)}$. Since L is finitely generated over $\mathbb{C}(t, s)$ and contained in $\overline{\mathbb{C}(t, s)}$ we have L finite extension of $\mathbb{C}(t, s)$. Let S' be a surface such that $\mathbb{C}(S')$ is isomorphic to L, respectively and S' maps finitely onto $\mathbb{P}^1 \times \mathbb{P}^1$. Then we have $A_{(t,s)}$ and $J(C_t) \times J(C_s)$ defined over L and we can spread $A_{(t,s)}$ and $J(C_t) \times J(C_s)$ over some Zariski open U in S'. Call these spreads as $\mathscr{A}$, $\mathscr{J}$. Then throwing out some more points from U we will get that the morphism from $\mathscr{A}$, $\mathscr{J}$ to U is a proper, submersion of smooth manifolds, if we view everything over $\mathbb{C}$(again here we use the non-canonical isomorphism $\overline{\mathbb{C}(t, s)} = \mathbb{C}$). Then by Ehressmann's theorem we have two fibrations $\mathscr{A} \to U$ and $\mathscr{J} \to U$. Since any fibrartion gives rise to a local system and hence a monodromy representation of the fundamental group of $\pi_1(U, t')$ on $H^{2d-1}(A_{(t,s)}, \mathbb{Q}), H^{2d'-1}(J(C_t) \times J(C_s), \mathbb{Q}) \cong H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$ where d, d' are dimensions of $A_{(t,s)}, J(C_t) \times J(C_s)$ (here we might have to replace (t, s) by (t', s'), but for very two general

points the fibers $A_{(t,s)}, A_{(t',s')}$ will be isomorphic so are $J(C_t) \times J(C_s)$ and $J(C'_t) \times J(C'_s)$). Now $\pi_1(U, t')$ is a finite index subgroup of G. We prove that $H = H^{2d-1}(A_{(t,s)}, \mathbb{Q})$ is a G-equivariant subspace of $H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$ (since $A_{(t,s)}$ is a sub-abelian variety of $J(C_t) \times J(C_s)$, $H^{2d-1}(A_{(t,s)}, \mathbb{Q})$ is a subspace of $H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$). Now G acts on $H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$ by the Picard-Lefschtez formula, that is

$$\gamma \times \gamma'.(\alpha + \beta) = \gamma.\alpha + \gamma'.\beta$$

which is equal to

$$\alpha - \langle \alpha, \delta_\gamma \rangle \delta_\gamma + \beta - \langle \beta, \delta_{\gamma'} \rangle \delta_{\gamma'} \ .$$

Now suppose that $\alpha + \beta$ belongs to H. We have to prove that for all $\gamma \times \gamma'$ in G, $\gamma \times \gamma'.\alpha + \beta$ belongs to H. Consider

$$(\gamma \times \gamma')^m(\alpha + \beta) = \alpha - m\langle \alpha, \delta_\gamma \rangle \delta_\gamma + \beta - m\langle \beta, \delta_{\gamma'} \rangle \delta_{\gamma'}$$

δ_γ is the vanishing cycles corresponding to γ. Since $(\gamma \times \gamma')^m$ is in $\pi_1(U, t')$ we have

$$(\gamma \times \gamma)^m(\alpha + \beta) - \alpha - \beta$$

is in H. That would mean that

$$m\langle \alpha, \delta_\gamma \rangle \delta_\gamma + m\langle \beta, \delta_{\gamma'} \rangle \delta_{\gamma'}$$

is in H, by applying the Picard Lefschetz once again we get that

$$(\gamma \times \gamma')^m(\alpha + \beta)$$

is in H. So H is G equivariant, hence it is either $\{0\}$ or $H^1(C_t, \mathbb{Q}), H^1(C_s, \mathbb{Q})$ or all of $H^1(C_t, \mathbb{Q}) \oplus H^1(C_s, \mathbb{Q})$. So the corresponding $A_{(t,s)}$ will either be zero or $J(C_t)$ or $J(C_s)$ or $J(C_t) \times J(C_s)$. □

This proves that if for one very general (t, s), $A_{(t,s)}$ is one of the above mentioned possibilities then for another very general (t', s'), $A_{(t',s')}$ will achieve the same possibility.

Theorem 3.2. *Suppose $A_0(S)$ is not isomorphic to the Albanese variety $Alb(S)$. Consider a net of Lefschetz pencils on S as before. Then for a very general (t, s), $A_{(t,s)}$ is actually $\{0\}$ or $J(C_t)$ or $J(C_s)$.*

Proof. This follows by analyzing the argument of theorem 19 in [BG]. □

3.3. **Self products of curves on self product of K3 surfaces.** Let us consider a $K3$ surface S and embed it into some $\mathbb{P}^N$. Let us take a Lefschetz pencil D on S, then for a general t in D we have C_t smooth and irreducible in S. So we have a closed embedding of $C_t \times C_t$ in $S \times S$. Then we consider the push-forward induced by this embedding, denote it by j_{t*} from $A_0(C_t \times C_t)$ to $A_0(S \times S)$. Since $A_0(C_t) \times A_0(C_t)$ maps surjectively onto $A_0(C_t \times C_t)$, we get that $A_0(C_t \times C_t)$ is weakly representable and hence $A_0(C_t \times C_t)$ is isomorphic to $Alb(C_t \times C_t)$, see [Vo][proof of theorem 10.11]. Then arguing as in proposition 6 in [BG] we get the following.

Proposition 3.4. *The kernel of j_{t*} is a countable union of translates of an abelian subvariety A_{0t} of the Albanese variety $Alb(C_t \times C_t)$.*

Proof. See proposition 6 in [BG]. □

A simple computation using Kunneth theorem and Lefschetz Hyperplane theorem shows that $H^7(S \times S, \mathbb{Q})$ is zero. Also $H^3(C_t \times C_t, \mathbb{Q}) = H^1(C_t, \mathbb{Q}) \oplus H^1(C_t, \mathbb{Q})$. This tells us that there is a natural action of $\pi_1(D \setminus \{0_1, \cdots, 0_m\}, t)$ on $H^3(C_t \times C_t, \mathbb{Q})$, which has the property that any $\pi_1(D \setminus \{0_1, \cdots, 0_m\}, t)$ equivariant subspace of it is either $\{0\}$ or $H^1(C_t, \mathbb{Q})$ or $H^3(C_t \times C_t, \mathbb{Q})$. Now we prove the following.

Theorem 3.5. *A_{0t} is either 0, or an abelian variety isogenous to $J(C_t)$ or it is all of $Alb(C_t \times C_t)$.*

Proof. Let us consider a t such that C_t is smooth. Consider the abelian variety A_{0t} and $Alb(C_t \times C_t)$. Consider the non-canonical isomorphism of $\overline{\mathbb{C}(x)}$ with $\mathbb{C}$ and view $A_{0t}, Alb(C_t \times C_t)$ as schemes over $\overline{\mathbb{C}(x)}$. Let L be the minimal field of definition of $A_{0t}, Alb(C_t \times C_t)$ in $\overline{\mathbb{C}(x)}$. Then L is a finite extension of $\mathbb{C}(x)$ and let C be a curve mapping finitely onto $\mathbb{P}^1$ and have function field L. Then we spread $A_{0t}, Alb(C_t \times C_t)$ over some Zariski open U in C. Denote the spreads by $\mathscr{A}_0, \mathscr{A}$ over U. Now throwing out some more points from U we can assume that $\mathscr{A}_0 \to U, \mathscr{A} \to U$ are proper submersions. Therefore the morphisms $\mathscr{A}_0 \to U, \mathscr{A} \to U$ are fibrartions by the Ehressmann's fibration theorem. So this will give us that, $H^{2d-1}(A_{0t}, \mathbb{Q}), H^{2g-1}(Alb(C_t \times C_t), \mathbb{Q})$ are $\pi_1(U, t')$ modules for some t' which maps to t. Here d, g are dimensions of $A_{0t}, Alb(C_t \times C_t)$. Now $H^{2g-1}(Alb(C_t \times C_t), \mathbb{Q})$ corresponds to the

Hodge structure $H^1(C_t,\mathbb{Q})\oplus H^1(C_t,\mathbb{Q})$ and since A_{0t} lies in $Alb(C_t\times C_t)$ we have that $H^{2d-1}(A_{0t},\mathbb{Q}) = H_t$ is inside $H^1(C_t,\mathbb{Q})\oplus H^1(C_t,\mathbb{Q})$. Now we prove that H_t is a $\pi_1(\mathbb{P}^1\setminus\{0_1,\cdots,0_m\},t)$-equivariant module. For that we have to prove that for a generator γ of the above group and for an α in H_t, $\gamma.\alpha$ is again in H_t. By the Picard Lefschetez formula we have

$$\gamma.\alpha = \alpha - \langle\alpha,\delta_\gamma\rangle\delta_\gamma$$

δ_γ is the vanishing cycle corresponding to γ. Since $\pi_1(U,t')$ is a finite index subgroup in $\pi_1(\mathbb{P}^1\setminus\{0_1,\cdots,0_m\},t)$, we get that there exists m such that γ^m belongs to $\pi_1(U,t')$. Then we get that

$$\gamma^m.\alpha - \alpha = m\langle\alpha,\delta_\gamma\rangle\delta_\gamma$$

is in H_t. Applying Picard-Lefschetz formula once again we get that $\gamma.\alpha$ is in H_t. This proves that H_t is $\pi_1(\mathbb{P}^1\setminus\{0_1,\cdots,0_m\},t)$-equivariant and hence we get that H_t is either $\{0\}$ or $H^1(C_t,\mathbb{Q})$ or $H^1(C_t,\mathbb{Q})\times H^1(C_t,\mathbb{Q})$. Consequently we get that A_{0t} is either $\{0\}$ or isogenous to $J(C_t)$ or $Alb(C_t\times C_t)$. □

Theorem 3.6. *For a very general t, the abelian variety A_{0t} is actually $\{0\}$ or isogenous to $J(C_t)$.*

Proof. Follows from theorem 19 in [BG]. □

3.7. **Rank one projective bundles on K3 surfaces and algebraic cycles.** . Consider a $K3$ surface S and a rank 2 vector bundle of S. Let us embed S inside some $\mathbb{P}^N$. Consider a smooth and irreducible hyperplane section C_t of S. Then we have the following commutative diagram.

$$\begin{array}{ccc} E_t & \longrightarrow & E \\ \downarrow & & \downarrow \\ C_t & \longrightarrow & S \end{array}$$

where E_t is the pullback of E by the closed embedding of C_t into S. Then this gives the following commutative diagram at the level of projective bundles.

$$\begin{array}{ccc} \mathbb{P}(E_t) & \longrightarrow & \mathbb{P}(E) \\ \downarrow & & \downarrow \\ C_t & \longrightarrow & S \end{array}$$

Then we can ask what is the kernel of the natural push-forward from $A_1(\mathbb{P}(E_t))$ to $A_1(\mathbb{P}(E))$. By the projective bundle formula we have the following that

$$A_1(\mathbb{P}(E_t)) \cong A_0(C_t), \quad A_1(\mathbb{P}(E)) \cong A_0(S)$$

then we have the following commutative diagram (it follows from the projective bundle formula that such a diagram is indeed commutative)

$$\begin{array}{ccc} A_1(\mathbb{P}(E_t)) & \longrightarrow & A_1(\mathbb{P}(E)) \\ \downarrow & & \downarrow \\ A_0(C_t) & \longrightarrow & A_0(S) \end{array}$$

Since the vertical arrows are isomorphisms we get that the kernel of

$$A_1(\mathbb{P}(E_t)) \to A_1(\mathbb{P}(E)) \ ,$$

is nothing but the kernel of

$$A_0(C_t) \to A_0(S) \ .$$

Now considering a Lefschetz pencil on S we can prove by theorem 19 in [BG] that for a very general t in the Lefschetz pencil, the kernel of the push-forward from $A_1(\mathbb{P}(E_t))$ to $A_1(\mathbb{P}(E))$ is countable.

3.8. **Branched covers of K3 surfaces.** Let S be a $K3$ surface and let $\widetilde{S}$ be a branched cover of S. Let us embed S into some $\mathbb{P}^N$. Then consider the smooth hyperplane sections of S inside $\mathbb{P}^N$. By Bertini's theorem a general hyperplane section C_t of S corresponds to a smooth, irreducible curve $\widetilde{C}_t$ inside $\widetilde{S}$. Now arguing as in theorem 6 in [BG] we can prove that the kernel of j_{t*} from $J(\widetilde{C}_t)$ to $A_0(S)$ is a countable union of translates of an abelian variety A_t of $J(\widetilde{C}_t)$. Now take a Lefschetz pencil D on S. We prove that for a general t in D, A_t is either $\{0\}$ or

$J(\widetilde{C}_t)$. Let us blow up the ramification locus of C_t and we can assume that $\widetilde{C}_t \to C_t$ is actually a covering. Now consider A_t inside $J(\widetilde{C}_t)$. Let us consider them over $\overline{\mathbb{C}(x)}$ via the isomorphism of $\overline{\mathbb{C}(x)}$ with $\mathbb{C}$. Let L be the minimal field of definition of $A_t, J(\widetilde{C}_t)$ and L is a finite extension of $\mathbb{C}(x)$. So let C be a curve such that it maps finitely onto $\mathbb{P}^1$ and the function field of C is L. Then spread $A_t, J(\widetilde{C}_t)$ over some Zariski open U in C. Then throwing out some more points of U, we can assume that we have two fibrations $\mathscr{A} \to U$ and $\mathscr{J} \to U$, whose generic fibers are A_t and $J(\widetilde{C}_t)$. This gives us that $\pi_1(U, t')$ acts on $H_t = H^{2d-1}(A_t, \mathbb{Q})$ and $H^{2g'-1}(J(\widetilde{C}_t), \mathbb{Q})$, here d, g' are dimensions of $A_t, J(\widetilde{C}_t)$. The later vector space is nothing but $H^1(\widetilde{C}_t, \mathbb{Q})$ and we have $H^1(C_t, \mathbb{Q})$ is embedded into it. So consider $H_t \cap H^1(C_t, \mathbb{Q}) = H'_t$. Now $\pi_1(U, t')$ is a finite index subgroup of $\pi_1(\mathbb{P}^1 \setminus \{0_1, \cdots, 0_m\}, t)$.

Now the standard Picard-Lefschetz formula argument and the fact that $\pi_1(U, t')$ is a finite index subgroup of $\pi_1(\mathbb{P}^1 \setminus \{0_1, \cdots, 0_m\}, t)$ we get that H'_t is a $\pi_1(\mathbb{P}^1 \setminus \{0_1, \cdots, 0_m\}, t)$ module embedded into $H^1(C_t, \mathbb{Q})$. Since the action of $\pi_1(\mathbb{P}^1 \setminus \{0_1, \cdots, 0_m\}, t)$ on $H^1(C_t, \mathbb{Q})$ is irreducible by theorem 3.27 in [Vo] we get that H'_t is either $\{0\}$ or all of $H^1(C_t, \mathbb{Q})$. So it would mean that there is an ablian subvariety B_t of A_t which is either $\{0\}$ or isogenous to $J(C_t)$.

Theorem 3.9. *The kernel of j_{t*} is a countable union of a translate of an abelian variety A_t inside $J(\widetilde{C}_t)$. Further this abelian variety A_t contains an abelian variety B_t, which is either zero or isogenous to $J(C_t)$.*

Now if B_t is isomorphic to $J(C_t)$, then we have A_t maps onto $J(C_t)$. That would mean that $J(C_t)$ maps to zero under the push-forward from $J(C_t)$ to $A_0(S)$. But for a very general t, the kernel of the push-forward from $J(C_t)$ to $A_0(S)$ is countable. Therefore either the very general member C_t of the Lefschetz pencil we started with is rational or B_t is 0.

References

[BG] K. Banerjee and V. Guletskii, *Rational equivalence for line configurations on cubic hypersurfaces in $\mathbb{P}^5$.*, `arXiv:1405.6430v1`, 2014.

[Voi] C.Voisin,*Symplectic invoultions of K3 surfaces act trivially on CH_0*, Documenta Mathematicae 17, 851-860, 2012.

[Fu] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 2. Springer-Verlag, Berlin, 1984.

[Ha] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.

[Vo] C. Voisin, *Complex algebraic geometry and Hodge theory II*, Cambridge University Press, Cambridge, 2003.

Indian Statistical Institute, Bangalore Center, Bangalore 560059
E-mail address: kalyanb_vs@isibang.ac.in